

# 40+ Doctors Tell UK Drug Regulators: Vaccinating Kids for COVID is ‘Irresponsible, Unethical and Unnecessary’

[40+ Doctors Tell UK Drug Regulators: Vaccinating Kids for COVID is ‘Irresponsible, Unethical and Unnecessary’](#)

*In an open letter to the Medicines and Healthcare Products Regulatory Agency, more than 40 doctors, medics and scientists in the UK say children are more vulnerable to the potential long-term effects of COVID vaccines.*

by [Children’s Health Defense Team](#), [The Defender](#)

June 1, 2021

A group of more than 40 doctors, medics and scientists is calling the UK government’s plan to vaccinate children for COVID “irresponsible, unethical and unnecessary.”

In an [open letter](#) addressed to the Medicines and Healthcare Products Regulatory Agency, the group said no one under 18 should be vaccinated for [COVID](#) because evidence shows the virus poses almost [no risk](#) to healthy children.

The letter was written in response to leaked [documents](#) from the UK National Health service detailing how UK health authorities plan to give COVID vaccines to children over 12 years old.

According to the documents, children as young as 5 years old could be vaccinated as early as July – a move the government

said may be necessary to keep COVID cases down as lockdowns lift across the country.

The authors of the letter said the [risk of death](#) from COVID in healthy children is 1 in 1.25 million. COVID vaccines, however, are linked to strokes due to cerebral venous thromboses in people under 40 – a [finding](#) that “led to the suspension of the [Oxford-AstraZeneca](#) children’s trial,” the authors said.

The letter states:

*“Children have a lifetime ahead of them, and their immunological and neurological systems are still in development, making them potentially more vulnerable to [adverse effects](#) than adults.”*

**Read “COVID-19 Child Vaccination: Safety and Ethical Concerns – An Open Letter From UK Doctors to Dr. June Raine, Chief Executive, MHRA”:**

We wish to notify you of our grave concerns regarding all proposals to administer COVID-19 vaccines to children. Recently [leaked Government documents](#) suggested that a COVID-19 vaccine rollout in children over 12 years old is already planned for September 2021, and the possibility of children as young as 5 years old being vaccinated in the summer in a worst-case scenario.

We have been deeply disturbed to hear several Government and SAGE representatives calling in the media for the COVID-19 vaccine rollout to be “[turning to children as fast as we can](#)”. [Teaching materials](#) circulated to London schools contain emotionally loaded questions and inaccuracies. In addition, there has been disturbing language used by teaching union leaders, implying that coercion of children to accept the COVID-19 vaccines through [peer pressure](#) in schools was to be encouraged, despite the fact that coercion to accept a medical

treatment is against UK and International Laws and Declarations. Rhetoric such as this is irresponsible and unethical, and encourages the public to demand the vaccination of minors with a product still at the research stage and about which no medium- or long-term effects are known, against a disease which presents no material risk to them. A summary of our reasons is given below and a more detailed fully referenced [explanation](#) is available.

## **Risks and benefits in medical treatments**

Vaccines, like any other medical treatment, come with varied risks and benefits. Therefore, we must consider each product, individually, on its merits, and specifically for which patients or sections of the population is the risk/benefit ratio acceptable. For COVID-19 vaccines, the potential benefits are clear for the elderly and vulnerable, however, for children, the balance of benefit and risk would be quite different. We are raising these concerns as part of an informed debate, which is a vital part of the proper, scientific process. We must ensure that there is no repeat of any past tragedies which have occurred especially when vaccines are rushed to market. For example, the swine flu vaccine, Pandemrix, rolled out following the pandemic of 2010, [resulted in over one thousand cases of narcolepsy](#), a devastating brain injury, in children and teenagers, before being withdrawn. Dengvaxia, a new vaccine against Dengue, was also rolled out to children ahead of the full trial outcomes, and [19 children died](#) of possible antibody dependent enhancement (ADE) before the vaccine was withdrawn. We must not risk a repeat of this with the COVID-19 vaccines, which would not only impact on the children and families affected, but would also have a hugely damaging effect on vaccination uptake in general.

No medical intervention should be introduced on a 'one size fits all' basis, but instead should be fully assessed for suitability according to the characteristics of the age cohort

and of the individuals concerned, weighing up the risk versus benefit profile for each cohort and the individuals within a group. This approach was outlined last October, by the head of the Government Vaccine Task Force, [Kate Bingham, who said](#) “We just need to vaccinate everyone at risk. There’s going to be no vaccination of people under 18. It’s an adult-only vaccine, for people over 50, focusing on health workers and care home workers and the vulnerable.”

## **Children do not need vaccination for their own protection**

Healthy children are at almost no risk from COVID-19, with [risk of death](#) as low as 1 in 2.5 million. No previously healthy child under the age of 15 died during the pandemic in the UK and admissions to hospital or intensive care are [exceedingly rare](#) with most children having no or very mild symptoms. Although Long-Covid has been cited as a reason for vaccinating children, there is little hard data. It appears [less common](#) and much [shorter-lived](#) than in adults and none of the vaccine trials have studied this outcome. The inflammatory condition, PIMS, was listed as a [potential adverse effect](#) in the Oxford AstraZeneca children’s trial. [Naturally acquired immunity](#) will give broader and better lasting immunity than vaccination. Indeed, [many children](#) will already be immune. Individual children at very high risk can already [receive vaccination](#) on compassionate grounds.

## **Children do not need vaccination to support herd immunity**

Already, [two thirds](#) of the adult population have received at least one dose of a COVID-19 vaccine. Models that assume vaccination of children is required to reach herd immunity have failed to account for the proportion who had immunity prior to March 2020 and those who have [acquired it naturally](#). Recent modelling suggested that the UK [had achieved](#) the required herd immunity threshold on 12 April 2021.

Children [do not transmit](#) SARS-CoV-2 as readily as adults, moreover adults living or working with young children are at lower risk of severe COVID-19. Schools [have not been shown](#) to be the focus on spread to the community, teachers have a lower risk of COVID-19 than other working age adults.

### **Short-term safety concerns**

As of 13th May, the MHRA22 has received a total of 224,544 adverse events, including 1,145 deaths in association with SARS-CoV-2 vaccines. Reports of strokes due to cerebral venous thrombosis were initially in low numbers but as awareness increased, many more reports led to the conclusion that AstraZeneca vaccine should not be used for adults under 40 years of age and this unpredicted finding has also led to the suspension of the Oxford AstraZeneca children's trial.

Similar events have been noted with Pfizer & Moderna vaccines on the US [adverse reporting system](#) (VAERS) and it is likely that this is a class effect related to production of spike protein. New UK [guidelines](#) on managing Vaccine-Induced Thrombotic Thrombocytopenia (VITT) include all COVID-19 vaccines in their advice. The possibility of further unexpected safety issues cannot be ruled out. In Israel, where the vaccines have been widely rolled out to young people and teenagers, the Pfizer vaccine has been linked to several [cases of myocarditis](#) in young men and concerns have been raised about [reports](#) of altered menstrual cycles and abnormal bleeding in young women following the vaccine.

Most concerning with regard to possible vaccination of children, is that there have now been a number of deaths associated with vaccination [reported to VAERS](#) in the US, despite the vaccines only being given to children within trials and a very recent rollout to 16-17 year olds.

### **Long-term safety concerns**

All Phase 3 COVID-19 vaccine trials are ongoing and not due to

conclude until late 2022/early 2023. The vaccines are, therefore, currently experimental with only limited short-term and no long-term adult safety data available. In addition, many are using a completely [new mRNA vaccine technology](#), which has never previously been approved for use in humans. The mRNA is effectively a pro-drug and it is not known how much spike protein any individual will produce. Potential late-onset effects can take months or years to become apparent. The limited children's trials undertaken to date are totally underpowered to rule out uncommon but severe side effects.

Children have a lifetime ahead of them, and their immunological and neurological systems are still in development, making them potentially more vulnerable to adverse effects than adults. A number of [specific concerns](#) have been raised already, including autoimmune disease and possible effects on placentation and fertility. A recently published paper raised the possibility that mRNA COVID-19 vaccines [could trigger](#) prion-based, neurodegenerative disease. All potential risks, known and unknown, must be balanced against risks of COVID-19 itself, so a very different benefit/risk balance will apply to children than to adults.

## **Conclusion**

There is important wisdom in the Hippocratic Oath which states, "First do no harm". All medical interventions carry a risk of harm, so we have a duty to act with caution and proportionality. This is particularly the case when considering mass intervention in a healthy population, in which situation there must be firm evidence of benefits far greater than harms. The current, available evidence clearly shows that the risk versus benefit calculation does NOT support administering rushed and experimental COVID-19 vaccines to children, who have virtually no risk from COVID-19, yet face known and unknown risks from the vaccines. The Declaration of the Rights of the Child [states that](#), "the child, by reason of his physical and mental immaturity, needs

special safeguards and care, including appropriate legal protection". As adults we have a duty of care to protect children from unnecessary and foreseeable harm.

We conclude that it is irresponsible, unethical and indeed, unnecessary, to include children under 18 years in the national COVID-19 vaccine rollout. Clinical trials in children also pose huge ethical dilemmas, in light of the lack of potential benefit to trial participants and the unknown risks. The end of the current Phase 3 trials should be awaited as well as several years of safety data in adults, to rule out, or quantify, all potential adverse effects.

We call upon our governments and the regulators not to repeat mistakes from history, and to reject the calls to vaccinate children against COVID-19. Extreme caution has been exercised over many aspects of the pandemic, but surely now is the most important time to exercise true caution – we must not be the generation of adults that, through unnecessary haste and fear, risks the health of children.

### **Signatories**

Dr Rosamond Jones, MD, FRCPCH, retired consultant paediatrician

Lord Moonie, MBChB, MRCPsych, MFCM, MSc, House of Lords, former parliamentary undersecretary of state 2001-2003, former consultant in Public Health Medicine

Prof Anthony Fryer, PhD, FRCPath, Professor of Clinical Biochemistry, Keele University

Professor Karol Sikora, MA, MBBChir, PhD, FRCR, FRCP, FFPM, Dean of Medicine, Buckingham

University, Professor of Oncology

Professor Angus Dalglish, MD, FRCP, FRACP, FRCPath, FMed Sci, Professor of Oncology, St Georges Hospital London

Professor Richard Ennos, MA, PhD. Honorary Professorial Fellow, University of Edinburgh

Professor Anthony J Brookes, Department of Genetics & Genome Biology, University of Leicester

Dr John A Lee, MBBS, PhD, FRCPath, retired Consultant Histopathologist, former Clinical Professor

of Pathology at Hull York Medical School

Dr Alan Mordue, MBChB, FFPH (ret). Retired Consultant in Public Health Medicine & Epidemiology

Dr Elizabeth Evans, MA, MBBS, DRCOG, retired doctor

Mr Malcolm Loudon, MB ChB, MD, FRCSEd, FRCS (Gen Surg). MIHM, VR. Consultant Surgeon

Dr Gerry Quinn, Microbiologist

Dr C Geoffrey Maidment, MD, FRCP, retired consultant physician

Dr K Singh, MBChB, MRCP, general practitioner

Dr Pauline Jones MB BS retired general practitioner

Dr Holly Young, BSc, MBChB, MRCP, Consultant physician, Croydon University Hospital

Dr David Critchley, BSc, PhD, 32 years in pharmaceutical R&D as a clinical research scientist.

Dr Padma Kanthan, MBBS, General practitioner

Dr Thomas Carnwath, MBBCh, MA, FRCPsych, FRCGP, consultant psychiatrist

Dr Sam McBride BSc(Hons) Medical Microbiology & Immunobiology, MBBCh BA0, MSc in Clinical

Gerontology, MRCP(UK), FRCEM, FRCP(Edinburgh). NHS Emergency



## Medicine & geriatrics

Dr Helen Westwood MBChB MRCP DCH DRCOG, general practitioner

Dr M A Bell, MBChB, MRCP(UK), FRCEM, Consultant in Emergency Medicine, UK

Mr Ian F Comaish, MA, BM BCh, FRCOphth, FRANZCO, Consultant ophthalmologist

Dr Jayne LM Donegan MBBS, DRCOG, DCH, DFFP, MRCP, general practitioner

Dr Dayal Mukherjee, MBBS MSc

Dr Clare Craig, BM,BCh, FRCPath, Pathologist

Mr C P Chilton, MBBS, FRCS, Consultant urologist emeritus

Dr Theresa Lawrie, MBBCh, PhD, Director, Evidence-Based Medicine Consultancy Ltd, Bath

Dr Jason Lester, MRCP, FRCR, Consultant Clinical Oncologist, Rutherford Cancer Centre, Newport

Dr Scott McLachan, FAIDH, MCSE, MCT, DSysEng, LL.M., MPhil., Postdoctoral researcher, Risk &

Information management Group

Michael Cockayne, MSc, PGDip, SCPHNOH, BA, RN, Occupational health practitioner

Dr John Flack, BPharm, PhD. Retired Director of Safety Evaluation at Beecham Pharmaceuticals

1980-1989 and Senior Vice-president for Drug Discovery 1990-92  
SmithKline Beecham

Dr Stephanie Williams, Dermatologist

Dr Greta Mushet, retired Consultant Psychiatrist in

Psychotherapy. MBChB, MRCPsych

Dr JE, MBChB, BSc, NHS hospital junior doctor

Mr Anthony Hinton, MBChB, FRCS, Consultant ENT surgeon, London

Dr Elizabeth Corcoran, MBBS, MRCPsych, Psychiatrist, Chair  
Down's Syndrome Research Foundation UK

Dr Alan Black, MB BS MSc DipPharmMed, retired pharmaceutical  
physician

Dr Christina Peers, MBBS, DRCOG, DFRH, FFRH, Consultant in  
Contraception & Reproductive Health

Dr Marco Chiesa, MD, FRCPsych, Consultant Psychiatrist &  
Visiting Professor, UCL

Elizabeth Burton, MB ChB, retired general practitioner

Noel Thomas, MA, MB ChB, DCH, DObstRCOG, DTM&H, MFHom, retired  
doctor

Malcolm Sadler, MBBS, FRCGP, retired general practitioner with  
37 years in Medical Practice

Dr Ian Bridges, MBBS, Retired general practitioner

© June 2021 Children's Health Defense, Inc. This work is reproduced and distributed with  
the permission of Children's Health Defense, Inc. Want to learn more from Children's  
Health Defense? [Sign up](#) for free news and updates from Robert F. Kennedy, Jr. and the  
Children's Health Defense. Your [donation](#) will help to support us in our efforts.

**[Connect with Children's Health Defense](#)**