

# Kids, Schools and COVID Tests – What Are the Risks?

## [Kids, Schools and COVID Tests – What Are the Risks?](#)

*Researchers warn of brain injury risks for millions of children who are now going to routinely undergo nasal swab tests administered by nonexperts – in some cases, by schoolchildren themselves.*

by [Alliance for Natural Health International](#)

March 22, 2021

Kids in the UK returned to school this month – the first step in a painfully slow easing of restrictions [tabled to last months](#). However, this is only on the grounds that children undergo regular [COVID-19](#) testing and follow mask mandates now extended into classrooms as well as common areas. All of which comes at a cost, both to the health and wellbeing of the children as well siphoning a [whopping £78 million](#) from taxpayers money.

But are these restrictions necessary? Research scientists from the Department of Infectious Disease at Imperial College, London, suggest not, because of the [extremely low SARS-CoV-2](#) transmission rates in schools.

### **Kids – fancy poking your brains?**

Some people are rightly concerned that the improper use of nasal swabs used in [PCR](#) or lateral flow testing may result in damage to the delicate membrane protecting the brain. But if you were to [believe the fact-checkers](#), you'd assume this is nothing more than a conspiracy theory. But as has been demonstrated in many other instances, the fact checkers have got it wrong again.

There is a risk – and that risk gets greater as more and more untrained people (now including children) administer their own swabs.

In October 2020, [a case report](#) was published in JAMA Otolaryngology-Head & Neck Surgery detailing the case of a woman whose brain membrane was pierced by a COVID test swab resulting in the leakage of cerebrospinal fluid. A second woman in the U.S. recently had [a similar experience](#). Assessing the use of nasal swabs, [researchers publishing in JAMA Otolaryngology-Head & Neck Surgery](#) issued warnings about the risks for the millions who are now routinely going to undergo tests, especially from ineptly administered nasal swabs.

And who's administering the nasal swabs in schools? The [schoolchildren themselves](#)! Let's hope that when kids do it wrong, it's because they don't go deep enough – not too deep. What no one seems to be being told is that the cribriform plate, the significant part that separates the brain from the nasal cavity is a delicate, soft, honey-comb or sieve-like structure that is thin and narrow with tiny perforations. This is why it's a perfect site for endoscopic transnasal skull-based surgery. But clearly it's proven nonsense to suggest someone inept at wielding a nasal swab can't push their swab into their brain.

### **Why lateral flow testing in schools is deeply misleading – and a waste of public money**

Nearly 57 million [lateral flow test \(LFT\) kits](#) have been delivered to schools in the UK.

Concerns have been leveled at their accuracy due to the number of [false positive results](#) they return. Conversely, there are also concerns over the high levels of false negatives, although this is less of an issue as true negative rates are far lower than is generally thought. Views around the use of mass testing to detect SARS-CoV-2 are mixed with concerns that mass testing [does not prevent transmission of the virus](#).

Confusion [reigns](#) in terms of positive tests. If a child tests positive at home and a subsequent PCR test comes back negative that child can return to school.

However, if a positive LFT test has been undertaken on school premises the pupil and their contacts [will have to self-isolate](#) whether or not they have a negative PCR test as the government considers the chances of the LFT test being a false positive as minimal.

We have previously discussed the problems with [PCR testing](#) and the significantly increased likelihood of a test returning a positive result when prevalence of the virus in the community is low. The same problem applies to lateral flow tests. To illustrate this we plugged in the sensitivity and specificity data (77.8% and 99.68% respectively) from the [Innova rapid flow test](#) into the [MedCalc Diagnostic test evaluation calculator](#). The aims were to calculate the positive predictive value (PPV) (i.e., the probability that SARS-CoV-2 is present when the test is positive) and the negative predictive value (NPV) (i.e., the probability that SARS-CoV-2 is not present when the test is negative) taking into account the low prevalence (amount) of infection in a given population.

This is important because, as we explained in our [‘Casedemic’ piece that critiqued PCR tests](#), Bayesian theory makes it essential that we take into account prevalence when considering the false positive and negative rates of any diagnostic test. A key fact that the UK’s health minister, Matt Hancock, just [can’t seem to wrap his head around](#) – or chooses not to.

We’ve used three prevalence rates: 0.14% (taken from [NHS Test & Trace data](#)), then halved that (0.07%) and doubled it (0.28%), to take into account variation, as prevalence is something of a moving feast.

At these prevalences, you’ll find much lower positive

predictive values (PPVs) than claims around accuracy being made by the government.

For the three prevalence rates, 0.07%, 0.14% and 0.28%, the chances of a positive test successfully indicating the presence of the infection is 15%, 25% and 41%, respectively. That's something school kids, parents and their teachers are just not being told.

The Royal Statistical Society's COVID-19 taskforce has [also raised concerns](#) over the risk of 'positive' tests when infection prevalence is low. Isn't it odd that no one seems to be listening? We'd ask again: where is the cost/benefit analysis showing that this massive purchase using taxpayers money of 57 million lateral flow test kits was justified for schools?

## **Spit or swab?**

Nose and throat swab testing can not only potentially harm the person being tested, it's also downright unpleasant. Is there a better way of testing? Actually there is, but we're not hearing much about it. It involves the plethora of saliva tests that [appear to be as effective](#) as the lateral flow tests – and far easier to administer. A [recent study](#) published in the New England Journal of Medicine found that a far higher number of saliva samples were positive for up to 10 days following diagnosis of COVID-19 compared to swab samples.

Saliva tests are becoming increasingly available, but have yet to be utilized by governments.

Here are some details:

### USA

- [SalivaDirect](#) – developed by Yale School of Public Health was [approved for use](#) by the U.S. Food & Drug Administration in August of last year.
- Scientists at Washington University have [announced the](#)

[development](#) of a saliva test they hope could be used at scale.

- [DxTerity](#) has added a saliva based SARS-CoV-2 test to its portfolio.

## UK

- [Covguard saliva tests](#) are available to UK citizens.
- Vatic Health is another UK based offering created by a team of scientists, engineers and designers.
- And if you're looking for saliva tests that also fulfill 'Fit to Fly' requirements then take a look at [Fitness Genes](#) and [Hydro-x](#).

## Australia

- A team at the [University of Technology, Sydney](#) has developed a very sensitive saliva test for SARS-CoV-2 antigens that can deliver results in under 15 minutes.

*Originally published by [Alliance for Natural Health International](#).*

Children's Health Defense, Inc. This work is reproduced and distributed with the permission of Children's Health Defense, Inc. Want to learn more from Children's Health Defense? [Sign up](#) for free news and updates from Robert F. Kennedy, Jr. and the Children's Health Defense. Your [donation](#) will help to support us in our efforts.